

GRACE-FO ML-calibrated magnetic field data

(<https://doi.org/10.5880/GFZ.2.3.2023.001>)

Kevin Styp-Rekowski^{1, 2, 3}, Ingo Michaelis², Monika Korte², Claudia Stolle⁴

1. *Distributed and Operating Systems, Technical University of Berlin, Ernst-Reuter-Platz 7, 10587 Berlin, Germany*
2. *GFZ German Research Centre for Geosciences, Helmholtz Centre Potsdam, Telegrafenberg, 14473 Potsdam, Germany*
3. *University of Rostock, Universitätsplatz 1, 18055 Rostock, Germany*
4. *Leibniz Institute of Atmospheric Physics at the University of Rostock, Schloßstraße 6, 18225 Kühlungsborn, Germany*

1. Licence

Creative Commons Attribution 4.0 International License (CC BY 4.0)



2. Citation

When using the data please cite:

Styp-Rekowski, K.; Michaelis, I.; Korte, M.; Stolle, C. (2023): GRACE-FO ML-calibrated magnetic field data. V. 0302. GFZ Data Services. <https://doi.org/10.5880/GFZ.2.3.2023.001>

The data are supplementary material to:

TBA

Table of contents

1. Licence	1
2. Citation	1
Table of contents.....	1
3. Data Description	2
3.1. Sampling method	2
3.2. Analytical procedure and data processing	2
4. File description	2
4.1. File naming convention	3
5. References	3

3. Data Description

The Gravity Recovery and Climate Experiment-Follow-On (GRACE-FO) satellite mission, consisting of two satellites, each carry a magnetometer as part of its attitude orbit control system (AOCS). After careful calibration, the data acquired through them can be used for scientific purposes by removing artificial disturbances from other satellite payload systems. This dataset is based on the dataset provided by Michaelis et al. (2022) and uses a similar format. The platform magnetometer data has been calibrated against CHAOS-7 magnetic field model predictions for core, crustal and large-scale magnetospheric field (Finlay et al., 2020) and is provided in the ‘chaos’ folder. The calibration results using a Machine Learning approach are provided in the ‘calcorr’ folder. Michaelis’ dataset can be used as an extension to this dataset for additional information, as they are connected using the same timestamps to match and relate the same data points. The exact approach based on Machine Learning is described in the referenced publication. Additionally, in the folder ‘fac’, field-aligned current derived from the magnetic field data are provided. The data is provided in NASA CDF format (<https://cdf.gsfc.nasa.gov/>) and accessible at: ftp://isdcdftp.gfz-potsdam.de/grace-fo/MAGNETIC_FIELD/ML/0302/ and further described in a README.

3.1. Sampling method

The data was recorded onboard the GRACE-FO satellites with varying time intervals of the different subsystems measuring. The magnetometer measurements (1s intervals) were aligned to match the closest position measurement (1s intervals) and interpolated accordingly. All other available data of potentially different intervals was interpolated and aligned to the same timestamps.

3.2. Analytical procedure and data processing

The data was calibrated using a Machine Learning approach involving Physics-informed Neural Networks, the whole method of calibration is described precisely in the referenced publication. The data was mainly processed for its calibration which yields a lower residual compared to a reference model than the uncalibrated data, more details about the many steps involved can be found in the referenced publication.

4. File description

The repository consists of three folders, calcorr, chaos and fac. The folders contain the CDF files of the calibration, whereby one file per day is available. The timestamps of every file in every folder matches with each other, thus allowing to combine the two datasets using the timestamp, if needed.

Hereby, the ‘chaos’ folder contains the CHAOS7 reference model values, each CDF file contains the position (encoded in latitude, longitude, and radius) as well as the prediction of the core, crustal and large-scale magnetospheric field in NEC frame as well as the combined value of the three contributions in NEC frame, as well as in satellite frame (MAG). In addition, the B-field estimates of the Average Magnetic field and Polar current System (AMPS) model are included.

The ‘calcorr’ folder contains the calibrated measurements of the platform magnetometer. Each CDF file contains the position (encoded in latitude, longitude, and radius) as well as the longitude and latitude according to the quasi-dipole coordinates. The calibrated measurements are provided in NEC and MAG frame. In addition, the quaternions are provided, which contain the rotation from the MAG to the NEC frame. Also, Flags are provided which are described in more detail in the referenced publication, ‘B_FLAG’ contains erroneous data, ‘KP_DST_FLAG’ shows magnetic activity as indicated by the Hp30 and Dst indices, ‘NaN_FLAG’ describes data points where NaNs have been automatically filled up with values.

The ‘FAC’ folder contains estimates for the Field-aligned currents as derived from the calibrated data in ‘calcorr’.

4.1. File naming convention

Exemplary: GF1_MAG_ACAL_CORR_ML_20091101T004915_20091101T235955_0302.cdf

With:

GF1:	GRACE-FO1 (satellite identifier, 1: First satellite)
MAG_ACAL_CORR_ML:	Magnetic field data, aligned, calibrated and corrected using ML (data category)
20180601T000000:	yyyymmddThhhhhh (time of first entry)
20180601T235959:	yyyymmddThhhhhh (time of last entry)
0302:	Version 0302 (version identifier)

5. References

Finlay, C. C., Kloss, C., Olsen, N., Hammer, M. D., Tøffner-Clausen, L., Grayver, A., & Kuvshinov, A. (2020). The CHAOS-7 geomagnetic field model and observed changes in the South Atlantic Anomaly. *Earth, Planets and Space*, 72(1), 156. <https://doi.org/10.1186/s40623-020-01252-9>

Michaelis, I., Stolle, C., & Rother, M. (2021). GRACE-FO calibrated and characterized magnetometer data (Version 0201) [dataset]. GFZ Data Services. <https://doi.org/10.5880/GFZ.2.3.2021.002>

NASA CDF Format: <https://cdf.gsfc.nasa.gov/>

TBA